

FIFRA Section 24(c) Special Local Need Label



Corteva Agriscience LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Strongarm®

EPA Reg. No. 62719-288

EPA SLN No. GA-040010

For Distribution and Use Only in the State of Georgia

Postemergence Application for Suppression of Tropical Spiderwort (*Commelina benghalensis*) Control in Peanuts

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read this SLN and the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this FIFRA 24(c) Special Local Need labeling is subject to all use precautions and limitations imposed by the label affixed to the container for this product.

Directions for Use

Apply Strongarm® herbicide in a broadcast spray when tropical spiderwort is in the 1 to 2 leaf stage of growth and actively growing. Applications made to larger weeds or to weeds under stress may result in unsatisfactory control. Optimum weed control will be obtained by applying Strongarm under favorable growing conditions (i.e., adequate soil moisture and temperature). Applications may occur from peanut cracking, to the initiation of flowering (bud stage), but no later than 30 days after planting.

Weeds Suppressed	Strongarm (oz/acre)
Tropical Spiderwort (<i>Commelina benghalensis</i>)	0.45

Surfactant Use: Applications of Strongarm must include a nonionic surfactant at 1 quart per 100 gallons (0.25% v/v). When mixing, add nonionic surfactant last.

Tank Mix Applications: Strongarm may be applied alone or in tank mix combination with other postemergence herbicides registered for use in peanuts. When applied in tank mix combination with other herbicides, follow applicable use instructions, including application rates, precautions and restrictions for each product used in the tank mixture, including use of adjuvants. The most restrictive limitations on each tank mix label must be followed.

Rotation to Cotton: Except for rotation to cotton, follow crop rotation intervals specified in the “Crop Rotation Intervals for Common Crops” section of the label booklet for Strongarm. For postemergence use in peanuts, the minimum interval for rotation to cotton is 18 months or a successful field bioassay. See instructions for Field Bioassay in the “Crop Rotation Interval for Common Crops” section of the label booklet for Strongarm.

Maximum Application Rates: In peanuts, do not apply more than 0.45 oz of Strongarm per acre per year in any combination of preplant incorporated, preplant surface, preemergence through cracking or postemergence applications.

Crop Tolerance: Strongarm may cause temporary yellowing or chlorosis of peanut foliage after application. These symptoms will vary in degree of severity, depending on environmental and crop growth factors (e.g., temperature and soil moisture). If these symptoms as described are not acceptable, do not use Strongarm as a postemergence application in peanuts.

™®Trademarks of Corteva Agriscience and its affiliated companies

R044-024

Approved: ___/___/___

Replaces R044-004 and R044-014